

METHOD FOR PRODUCING HIGHLY FUNCTIONAL, HYPERBRANCHED POLYESTERS

Publication number: WO03093343

Publication date: 2003-11-13

Inventor: WAGNER EVA (DE); BRUCHMANN BERND (DE);
KELLER PETER (DE)

Applicant: BASF AG (DE); WAGNER EVA (DE); BRUCHMANN
BERND (DE); KELLER PETER (DE)

Classification:






- international: C08G63/12; C08G63/81; C08G63/82; C08G63/00;
(IPC1-7); C08G63/02; C08G63/12; C08G63/40;
C08G63/60; C08G63/80; C08G63/87

- European: C08G63/12; C08G63/81

Application number: WO2003EP04121 20030422






Priority number(s): DE20021019508 20020430

Also published as:

 EP1501882 (A1)
 US2005185177 (A1)
 EP1501882 (A0)
 DE10219508 (A1)
 CN1649931 (A)

more >>

Cited documents:

 US4749728
 EP0680981
 US5663281
 US4668763
 JP63172727

more >>

Report a data error here

Abstract of WO03093343

The invention relates to a method for producing highly functional, hyperbranched polyesters. Said method is characterised in that (a) one or more dicarboxylic acids or one or more derivatives thereof is/are reacted with one or more at least trifunctional alcohols or (b) one or more tricarboxylic acids or higher polycarboxylic acids or one or more derivatives thereof is/are reacted with one or more diols in the presence of a solvent and optionally in the presence of an inorganic, organometallic or low-molecular organic catalyst.

Data supplied from the **esp@cenet** database - Worldwide